

Paleomagnetism and the Stability of Dharwar Craton, India

A.F. Pivarunas¹, J.G. Meert¹

¹University of Florida, Department of Geological Sciences, 241 Williamson Hall, Gainesville, FL 32611, afp5@ufl.edu

Abstract

The prominent bend of Dharwar Craton, India was recently suggested to post-date the ‘stabilization’ phase of the craton via a $\sim 30^\circ$, anticlockwise, vertical-axis intracratonic rotation. The trends of mafic dyke swarms were used as the prime piece of evidence to make this case. However, the trends of dyke swarms may not be the best method of assessing intracratonic rotation. We examine the large paleomagnetic database of Dharwar craton as an independent method to assess the rotation. We find that multiple early Paleoproterozoic dyke swarms (the 2367, 2216, and 2082 Ma generations) preserve a seemingly systematic paleomagnetic directional offset. The mean paleomagnetically permissible rotation is $-19^\circ \pm 17^\circ$. The large, Hampi dyke swarm, emplaced ~ 1888 Ma does not show the same offset, which suggests no relative disturbance of the cratonic nuclei took place after this time. Therefore, the 2082-1888 Ma interval brackets the window for any deformation and tectonism required to bend the Dharwar craton by about $\sim 20^\circ$. Unfortunately, the Dharwar craton is tectonically quiescent during this time period which is hard to reconcile with a craton-scale deformational event. We argue that the ‘observed’ rotation may result from biases associated with how trends of dykes and their magnetic directions are compiled when age data are limited.

Introduction

Lithospheric mobility on geologic timescales results in the interaction of lithospheric plates. Paleogeographic research strives to meticulously document the movement and interaction of the lithospheric plates over geologic time. Although the interactions of plates, and therefore orogenic effects, are principally focused along their margins, the interiors of plates do not escape deformation. Around 35 cratons have survived since the Archean (the so-called ‘puzzle pieces’ of Bleeker, 2003) and these are the relative pieces to consider when assessing intraplate, or rather intracratonic, deformation over the long term. Various datasets can be used to define intracratonic deformation: the coherence of ancient sedimentary basins, metamorphic zones, or the geometric pattern of linear features such as dyke swarms and structural trends. The best method, however, is paleomagnetism. Within the powerful, simple framework of the geocentric axial dipole hypothesis (GAD; Meert, 2014), paleomagnetic results from across a craton can be examined for consistency. Craton-wide uniformity of paleomagnetic results would support hypotheses that involve little to no intracratonic deformation. This approach is exemplified in the comprehensive paleomagnetic, geometric, and tectonic examination of the Paleoproterozoic Superior craton (Evans and

Halls, 2010), which found that the eastern half of the craton is best restored to the western half with a $\sim 14^\circ$ anticlockwise rotation. A necessary component to this style of analysis is a detailed paleomagnetic record from the craton in question.




Figure 1: Map of Dharwar craton showing the four major mafic dyke swarms that provided the paleomagnetic data used to test intracratonic rotation hypotheses.

Dharwar Craton is a large and well-studied block of ancient Archean crust that is exposed across much of Southern India (Naqvi and Rogers, 1987; Ramakrishnan and Vaidyanadhan, 2008; Meert et al., 2010). The craton hosts numerous swarms of mafic dykes of multiple geometrical trends which have been separated into chronological generations based on decades of concerted paleomagnetic and geochronologic study (Fig. 1). Recent summations of dyke ages and relationships delineate over 7 different emplacement events within the craton (Samal et al., 2015; Söderlund et al., 2019; Samal et al., 2019).

A distinct feature of Dharwar craton is a curvilinear structural grain which bends from south to north through the craton (Fig. 1; Naqvi and Rogers, 1987). This curvilinear trend is recognizable in three features: the Chitradurga Fault (shear zone), the Closepet Granite, and the Andhra Karnataka Long Dyke (AKLD). There are two end-member explanations for this: either this a primary (i.e. pre-stabilization, Archean) or secondary (i.e. post-stabilization, Proterozoic) feature. The possibility that Dharwar craton experienced an intracratonic rotation to produce this arcuate shape as a secondary feature was raised (Kumar et al., 2015) and dismissed (Nagaraju et al., 2018) in specific reference to the AKLD, a conspicuously curved 2216 Ma dyke which stretches >400 km across the craton from north to south. Recent work again

raised the issue of a late (i.e. Paleoproterozoic) relative rotation between north and south Dharwar craton (Söderlund et al., 2019). Söderlund et al. (2019) propose $\sim 30^\circ$ of anticlockwise vertical-axis rotation. According to their model, the correction transforms currently radiating 2367-2180 Ma dyke swarms into linear swarms. This approach relies on the assumption of original linearity of the dyke swarms.

In this paper, we examine the paleomagnetic data available from four major dyke swarms within Dharwar craton – the 2367, 2216, 2080 and 1885 Ma swarms – and their implications for relative intracratonic rotation. We undertake this analysis because we believe that paleomagnetic data provide a more robust test for rotation than does assuming an initial geometry of emplacement.

Assessing vertical axis rotation: direction-space and pole-space

The paleomagnetic database of Dharwar craton is now well-established, with robust results from mafic dykes in the north and south sectors of Dharwar craton (Fig. 1). We will not discuss the paleomagnetic data in detail here (for a discussion of the key paleomagnetic poles refer to Meert et al., in press). We gathered site-level paleomagnetic data from the four major dyke swarms with large enough datasets to be useful and analyzed them using Python software with a specialized paleomagnetic package (Tauxe et al., 2016). Paleomagnetic directional data (declination and inclination) are ideal for assessing vertical-axis rotation of the craton (see Butler, 1992). This is achieved simply by comparing the observed paleomagnetic declination in a given region of questionable tectonic stability with the expected paleomagnetic declination from a stable reference area.

$$R = D_O - D_X$$

The difference between the declinations (R) is the amount of relative rotation. For our purposes, we compare the paleomagnetic declination of north versus south Dharwar craton. The southern sector of the craton ($<14^\circ$ N) serves as our ‘stable’ reference (Fig. 2,3).

Uncertainty on estimates of rotation (ΔR) is a function of uncertainty of paleomagnetic declination (ΔD_O for the ‘observed’ direction and ΔD_X for the ‘expected direction’) along with a correction factor of 0.8 (Demarest, 1983). Directional data do not conform with a Fisher distribution (Tauxe and Kent, 2004; Deenen et al., 2011), and thus Fisher statistics are not appropriate for assessing the uncertainty on purely directional data. However, the use of A95, as calculated from the distribution of virtual geomagnetic poles (VGPs), which do conform to Fisherian statistics, enables us to use the Demarest (1983) uncertainty calculation method more appropriately. The main upshot is a slightly higher uncertainty on the paleomagnetic declinations and thus a larger interval where rotations are not statistically definable. If the amount of rotation (R) does not exceed the uncertainty on its estimate (ΔR), the directions compared are technically concordant even if R is non-zero. That is, the declination difference is paleomagnetically indistinguishable. If instead the defined rotation does exceed uncertainty, the directions are discordant, and a paleomagnetically definable rotation is suspected.

Paleomagnetic site-level data can also be used to calculate a paleomagnetic pole, which is generally assumed to represent the ancient position of the Earth’s geographic pole with respect to a given area, subject to typical paleomagnetic uncertainties (Butler, 1992). We calculated paleomagnetic poles as the mean of virtual geographic poles (VGPs) calculated from each site mean (Supplementary Material). The paleomagnetic poles from north and south Dharwar craton are then compared in pole-space both before and after Euler rotations (Fig. 4). The hypothesized vertical-axis rotation between the north and south of the craton is best modelled as a Euler pole within the craton itself. We placed the pole at the hypothesized axis of rotation (Euler pole at 15° N, 75° E) of Söderlund et al. (2019) and examined both the amount of rotation hypothesized in their paper along with the rotation implied by paleomagnetic data (Fig. 4).

A summary of the paleomagnetic parameters of the total site-level datasets and calculations associated with this paper is attached as a Python notebook, along with the data used in this analysis (Supplementary Material).

Rotation Evidence from Paleomagnetism




Figure 2: Stereoplots of mean directional disparity among the four dyke swarms examined in this study. Black circles with confidence intervals represent the directional mean from north (triangle) and south (square), while the faded circles (blue=north and orange=south) represent site-level paleomagnetic data. Difference in declination between means are shown as lines radiating from the center of the plots.

The 2367 Ma Bangalore-Karimnagar dyke swarm is widespread throughout the entire Dharwar craton, from the Southern Granulite Terrane (SGT) (Fig. 1; Halls et al., 2007; Dash et al., 2013; Belica et al., 2014; Pivarunas et al., 2019), through the rest of the craton (Kumar et al., 2012a; Söderlund et al., 2019) all the way into Bastar craton to the north (Liao et al., 2019). The trends of these dykes generally fan from NW-SE in the south to NE-SW in the north. Thus, it has been proposed that these dykes are part of a radiating swarm with a center to the present geographic west of Dharwar craton (Halls et al., 2007). In contrast, the recent hypothesis of intracratonic rotation holds that this swarm was roughly linear, and that the fanning of these dykes is a secondary feature (Söderlund et al., 2019).

Name	N	D (°)	I (°)	k	a95 (°)	PLat (°N)	PLon (°E)	K	A95 (°)	R (°)	ΔR (°)
2367N	26	76.6	-76.9	60.8	3.7	8.2	54.2	18.9	6.7	-23.0	24.3
2367S	57	99.6	-83.5	41.3	3.0	14.3	65.1	12.4	5.6		
2216N	13	233.8	-62.1	33.2	7.3	-36.3	302.9	16.9	10.4	-15.9	15.0
2216S	12	249.7	-58.5	37.9	7.1	-23.8	309.3	22.9	9.3		
2082N*	27	43.7	-1.1	28.2	5.3	43.5	186.5	37.2	4.6	-17.7	7.1
2082S*	7	61.4	1.3	32.9	10.7	27.8	174.7	64.4	7.6		
1888N	39	308.7	-3.8	13.1	6.6	36.2	334.1	19.5	5.3	4.1	7.8
1888S	23	304.7	-9.2	9.8	10.2	31.9	333.6	14.7	8.2		

Notes: Name=age of dyke swarm, N indicates upper craton, S indicates southern craton; N=site-level data incorporated into mean directions; Dec = paleomagnetic declination, Inc = paleomagnetic inclination, , k = kappa precision parameter (Fisher, 1953); a95 = cone of 95% confidence about the mean direction, PLat = virtual geomagnetic pole latitude, PLon = virtual geomagnetic pole longitude (these last two calculated from directional data and site location); K = kappa precision parameter (Fisher, 1953); A95 = circle of 95% confidence about the paleomagnetic pole; R=paleomagnetic rotation seen in directional space; ΔR=95% confidence limits on paleomagnetic rotation (Demarest, 1983).




Figure 3: Amount of rotation defined by the four major dyke swarms. The three early Paleoproterozoic dyke swarms indicate anti-clockwise rotation, however, by 1885, concordant paleomagnetic results indicate that intracratonic rotation had ceased.

The total amount of rotation we can observe from paleomagnetic data is calculated as the difference between the mean declination in the north and the mean declination in the south. For the 2367 Ma dykes (N=83), we find that $R = -23.0^\circ \pm 24.3^\circ$ when separating north and south Dharwar results at 14° N (Table 1; Fig. 2,3). Although the sense of rotation is anti-clockwise, the magnitude of the uncertainty exceeds the rotation and therefore the north and south Dharwar craton paleomagnetic data at 2367 Ma are not statistically discordant.

A 2216 Ma subswarm of dykes is a prominent member of ca. 2.2 Ga magmatism stretching across the craton (Kumar et al., 2012a; Nagaraju et al., 2018a; Nagaraju et al., 2018b). Other swarms slightly post-dating 2216 Ma may represent the same event; however, due to the absolute rotation of Dharwar craton as a whole during this time interval, we only examine the precisely-dated 2216 Ma subswarm. For the 2216 Ma dykes (N=25), we find that $R = -15.9^\circ \pm 15.0^\circ$ (Table 1; Fig. 2,3). In this case, the magnitude of rotation just rises above the level of uncertainty such that the paleomagnetic data from 2216 Ma are discordant. Detailed sampling of the AKLD, however, fails to show the expected local trend/declination relationship implied by intracratonic rotation post-emplacement (Nagaraju et al., 2018a).

The 2082 Ma dyke swarm mainly has been found to the north and west of Cuddapah Basin (Kumar et al., 2015). For this reason, a boundary placed at 14° N results in no available paleomagnetic data from the ‘southern’ part of the craton. Thus, we consider a boundary at 15° N for this time period. For the 2082 Ma dykes (N=34), we find that $R = -17.7^\circ \pm 7.1^\circ$ (Table 1; Fig. 2,3). This amount of rotation between north and south Dharwar craton reaches ‘discordant’ status. The trends of the 2082 Ma dykes, it should be emphasized, do not help define the hypothesized rotation between north and south Dharwar (Söderlund et al., 2019), this is only seen in the paleomagnetic data presented here.

A large igneous province (LIP) principally manifesting as a dyke swarm (sills/traps in the Cuddapah Basin) stretches across Dharwar craton and the neighboring Bastar craton to the north at ~1888 Ma (French et al., 2008). A total of 54 dykes/sills were collected from this LIP since the inception of paleomagnetic studies in Dharwar craton (Belica et al., 2014; Meert et al., in press). For the 1888 Ma LIP, we find that $R = +4.1^\circ \pm 7.8^\circ$ (Table 1; Fig. 2,3). No discordance of declination is definable between the upper and lower reaches of the craton. Thus, on the basis of paleomagnetism, no intracratonic deformation is indicated after 1888 Ma.

We take the mean (\bar{R}) of the 2367, 2216, and 2082 Ma paleomagnetically defined rotations to provide a single number of the amount of intracratonic rotation permissible from analysis of paleomagnetic results. Given the high variance of the calculated confidence intervals for two of three estimates of rotation, we present the pooled uncertainty ($\Delta\bar{R}$) as follows in order to account for the uncertainty on the estimates:

$$\Delta\bar{R} = \sqrt{\frac{\sum \Delta R_i^2}{N}}$$

$$\bar{R} \pm \Delta\bar{R} = -18.9^\circ \pm 16.9^\circ$$

This range encompasses, within confidence limits, the previous estimate for N-S intracratonic anticlockwise rotation of ~25° (Kumar et al., 2012, cf. Nagaraju et al., 2018a), and the 30° proposed by Söderlund et al., (2019). It is also compatible with negligible rotation.




Figure 4: Pole-space comparison of results from north and south Dharwar craton (squares with blue A95 are from north and circles with orange A95 are from south) in (a) present-day arrangement showing the relative separation in pole space and (b) 'corrected' for anticlockwise rotation using an Euler pole of [15, 75, -17.7]. The fit is quite noticeably improved for 2216 Ma and 2080 Ma results; however, the 2367 results do not improve significantly. Red

squares indicate the illustration of a 30° clockwise correction as indicated by Söderlund; this approach does not fit the paleomagnetic data as well.

For comparison in pole-space, our vertical-axis correction (clockwise rotation) is derived from the 2082 Ma dykes ($R = -17.7^\circ$) since this is the datum with the most reasonable confidence limits. The present-day offset between north and south poles is noticeable in mean pole positions (Fig. 4a). When corrected with 17.7° of clockwise rotation, the improved fit of the 2216 and 2082 Ma paleomagnetic poles is immediately apparent (Fig. 4b). We also used a 30° clockwise rotation for comparison that ‘overcorrects’ the paleomagnetic data. The 2367 Ma results do not show an improved fit in either case, perhaps because of the proximity of the poles to the axis of rotation.

Discussion

In our attempt to harmonize the paleomagnetic data and age/dyke trends with the structural grain of the Dharwar craton, we examine potential issues with our analysis and that of Söderlund et al. (2019).

Given the rather large confidence intervals on the paleomagnetic rotation data, particularly for the high-latitude results which characterize Dharwar craton in the early Paleoproterozoic, they are at first glance compatible with the hypothesis of intracratonic rotation. At 2082 Ma, particularly, there is statistically unambiguous paleomagnetic evidence of rotation although the locus for that analysis is different than that used for the older dykes. At 1888 Ma, there is similarly unambiguous paleomagnetic evidence of no rotation. Thus, any large-scale intracratonic deformation of Dharwar craton pre-dates 1888 Ma. But what are the older data indicating?

Positioning the location of the rotation axis is of key importance. For example, if we adjust the boundary for the 2367 Ma dykes to 15° N, there are fewer dykes in the northern sector (only 9 of 83 units), and $R = +9.9^\circ \pm 27.0^\circ$. This is the opposite sense of rotation from what is ‘expected’ and disagrees with results using the 14° N boundary; the statistical threshold for discordance is not reached in either case. The same is true for the 2216 Ma dykes. If we adjust the boundary to 15° N, our calculation changes to $R = +20^\circ \pm 41.1^\circ$ which is also the opposite sense of rotation. Since this situation occurs for both the 2367 Ma and 2216 Ma results at the 15°N sector boundary, this may be an artifact of comparing steep-intermediate inclination directional means from drastically different sample sizes. For 2082 Ma dykes, changing the boundary is a fruitless exercise given that no paleomagnetic results for 2082 Ma dykes exist below 14° N. This exercise in boundary condition adjustment should give one pause before unequivocally accepting the paleomagnetic data as strong evidence for rotation. For 1888 Ma dykes, we find $R = -0.4^\circ \pm 7.3^\circ$, which is similar to a 14°N boundary in both concordance of directional data and indication of no rotation; this is expected given that the datasets have comparable numbers for both cases. Thus, the only dataset that is not strongly influenced by boundary conditions is also the dataset which illustrates a *lack* of rotation.

We also should critically examine the uncritical use of dyke trends as an analogue for age. The Söderlund et al. (2019) hypothesis of intracratonic rotation relies on empirically unsupported assumptions regarding trends of dykes throughout Dharwar craton. For example, 2367 Ma dykes from southernmost part of the craton (Halls et al., 2007; Dash et al., 2013; Belica et al., 2014; Pivarunas et al., 2019), do not show ‘improved’ linearity after rotating northern trends clockwise by 30° and were simply ignored in the prior analysis (see Fig. 5 of Söderlund et al., 2019). The NW-SE trend we see in the northern SGT (which is simply an extension of Dharwar craton) remains unchanged given an axis of rotation at 15° N, 75° E. One can always rotate a portion of a radiating dyke swarm back to linearity with another portion through an appropriate angle of rotation. For example, if the 2367 Ma dyke swarms represent a giant-radiating swarm, only ~60° (the south-to-north change in dyke trends) of the total radiation is present. Thus, arguing that central and northern Dharwar craton dykes become approximately linear after ~30° of relative rotation is still consistent with a radiating dyke swarm, and is not *prima facie* evidence for either original linearity or subsequent deformation.

As an example of trend-age problems, India hosts the spectacular Deccan large igneous province (LIP) – a comparatively modern analogue – where the lack of a preferred orientation of Deccan feeder dykes indicates that we should not necessarily expect a preferred trend-age relationship for Proterozoic dyke swarms (Vanderkluyesen et al., 2011). This is abundantly clear in India from the coincidence of 2367 Ma and 1888 Ma trends (Samal et al., 2015). It is likely that both dyke swarms exploited preferred zones of weakness in the crust (Pivarunas et al., 2019). The pulsed generations of Proterozoic dykes could either represent the remnants of plumbing systems of older LIPs or be related to extensional tectonics (Ernst, 2014). Similarly, the ~1.2 Ga Harohalli alkaline dykes (Pradhan et al., 2009) share a general N-S trend with many 2.2 Ga mafic dykes (Kumar et al., 2012; Nagaraju et al., 2018a; Nagaraju et al., 2018b). The 2080 Ma dykes from around Cuddapah Basin have a generalized radiating structure, which does not improve on ‘correcting’ the hypothesized northerly deformation (Söderlund et al., 2019). Untangling the secondary effects of regional deformation on the basis of dyke swarms must take into account deviations from ideal ‘one-trend-one-age’ behavior.

Furthermore, it is important to remember that most mafic dykes lack radiometric ages. A common practice is to group directional data on the basis of comparison to dykes that are well-dated *and* have paleomagnetic data. Occasionally, the dykes are also correlated on the basis of geochemical similarities (see, for example Liao et al., 2019) or trend data without any reference to paleomagnetic data. This does result in somewhat circular criteria for including dykes in a given dataset, with directional comparison likely the most inclusive (not always a good thing!) metric. Directional comparison likely results in artificial tightening of directional groupings to some degree, but also may result in an artificial loosening of groupings – and thus declination differences. On the other hand, relying solely on trends of dated dykes,

or geochemical similarities can also create artificial ‘lumping’. This is well-illustrated by the southernmost 2367 Ma dykes in Dharwar craton; they either help define a radiating swarm, or else are conjugate to an otherwise linear swarm. In neither case do the 2367 Ma dykes all follow a single trend. It would be ideal, albeit impractical, to date every single paleomagnetically sampled dyke. We simply stress caution in the uncritical use of these approaches and note that the apparent rotations in the Dharwar craton should be viewed with caution.

Our final point relates to tectonic matters, the *corpus delicti* which should accompany intracratonic rotation. Thus, we emphasize that the most important results from this analysis are the ‘positive’ test of rotation at 2082 Ma, and the ‘negative’ test of rotation at 1888 Ma. This provides a window of time in which to search for evidence of either oroclinal bending or protracted shearing (Söderlund et al., 2019) in the Dharwar craton. In other well-documented examples of intracratonic deformation, the zone of accommodation is recognizable and falls within the proper temporal framework (Li and Evans, 2014; Evans and Halls, 2010). Thus, for Dharwar Craton, a simple question arises. *Where is the deformation?*

The Dharwar Craton is tectonically quiescent during the 2100-1900 Ma interval, which is precisely when our results indicate that deformation should occur. Any event, either brief or protracted, that produced a $\sim 20^\circ$ bend of the entire craton should leave some mark in the tectonic record. Thus, we have an apparent solution (paleomagnetically defined rotation) in search of a deformational accommodation. One option is that 2367 Ma and 2216 Ma dykes simply each produce a similar apparent anticlockwise rotation given the scatter inherent in the averaging of paleosecular variation at relatively high latitudes. In this case, the 2082 Ma rotation – which is statistically better defined – may have resulted from more localized accommodation around the Cuddapah Basin. The pole-space results are more intriguing, given that both 2216 Ma and 2082 Ma paleomagnetic poles show improved fit after the northerly results are corrected for the proposed rotation (Fig. 4). This gives us higher confidence in the 2216 Ma results, which only just exceed uncertainty in directional-space. We note that the 2216 Ma declination difference does not exceed uncertainty ($14.8^\circ \pm 17.1^\circ$) if a more stringent compilation (N=21) of dykes is used (Meert et al., in press).

The Neoproterozoic 40° clockwise rotation of the South and West Australia cratons (SAC and WAC) with respect to North Australia craton (NAC) provides an excellent example of harmony between tectonic and paleomagnetic data (Li and Evans, 2011). A cross-continental megashear zone is visible within the Neoproterozoic Paterson and Petermann orogenies (Myers et al., 1996). In this case, there is both a clear post-rotation fit from the paleomagnetic data and a clear zone of deformation accommodating the rotation. Another example of Proterozoic cratonic rotation is within Superior Craton (Halls and Evans, 2010). This rotation, although relatively ‘small’ ($\sim 14^\circ$), is accommodated by a major deformational area, the Kapuskasing Structural Zone (KSZ).

The major tectonized regions of the Indian shield that may be relevant to Dharwar craton rotation include the Central Indian Tectonic Zone (CITZ) and the Eastern Ghats Province (EGP). The CITZ suffered cryptic deformation at around 2.5 Ga in the Sausar Mobile Belt (Stein, 2004, 2006), but major tectonic activity took place between 1.6–1.5 Ga and again during the assembly of proto-India around the Mesoproterozoic-Neoproterozoic transition (Bhowmik et al., 2019). Deformation in the EGP is broadly coeval with that in the CITZ (Dobmeier and Raith, 2003). Thus, major Proterozoic deformation in both of these provinces post-dates the paleomagnetically defined hard lower limit on rotation in Dharwar craton (1888 Ma).

There have been multiple suggestions that the structural grain of Dharwar craton pre-dates the Proterozoic. Chadwick et al. (2000) posited a plate-tectonic paradigm for the assembly of Dharwar craton and credit a Neoproterozoic collision between the eastern and western Dharwar craton with creating the present curved structural grain. Chardon et al. (2008), disagreed with the mechanism and some interpretations of previous workers, yet still favored late Archean timing for development of the Dharwar craton crustal architecture. These scenarios are plausible hypotheses for the formation of the curvilinear structural grain that do not require major post-stabilization intracratonic rotation.

Conclusions

The trends of dykes provide insufficient evidence for diagnosing intracratonic rotation. Paleomagnetic studies are more robust because any rotation should be apparent in the data. There is limited evidence for an anti-clockwise rotation of the northern Dharwar craton based on paleomagnetic results between 2367 and 2082 Ma. The offset is dependent on the areal division of Dharwar craton into ‘north’ and ‘south’ sectors. Depending on how the boundary is drawn, the datasets show either limited evidence for anti-clockwise rotation or the opposite sense of rotation (clockwise). Rigorously honoring statistical constraints, the best evidence via paleomagnetic results is from the 2082 Ma dykes and suggests $\sim 17^\circ$ of counterclockwise rotation. By 1888 Ma, paleomagnetic results are concordant. Therefore, any event or long-term tectonic effect suggested as the ‘prime rotator’ of Dharwar craton is constrained to post-date 2082 Ma and pre-date 1888 Ma. The lack of any reasonable driver for craton-wide deformation during this interval necessitates alternate hypotheses for the paleomagnetic data. We believe the simplest explanation is that the structural grain of Dharwar craton pre-dates emplacement of the dykes and is a relic of collision between the EDC and WDC during the Archean. We believe the paleomagnetic data certainly preclude significant deformation of the Dharwar Craton in the Mesoproterozoic to Neoproterozoic.

Acknowledgements

This work was carried out while AFP was a Visiting Assistant Professor position at the University of Florida and benefited from support from the Geological Sciences department at the University of Florida.

The authors would like to thank the paleomagnetism group in the Neil Opdyke Laboratory at the University of Florida for providing insightful observations which greatly improved the manuscript. This material is based upon work supported by the National Science Foundation under Grant No. EAR18-0693 to JGM.

Data Availability

Excel files of site-level paleomagnetic information used in this analysis is provided, along with a Jupyter notebook that provides reproduction of analyses reported in this paper. If there are any questions related to the use of the notebook, please direct them to AFP, the corresponding author.

References

- Babu, N. R., Venkateshwarlu, M., Shankar, R., Nagaraju, E., & Parashuramulu, V., 2018. New paleomagnetic results on 2367 Ma Dharwar giant dyke swarm, Dharwar craton, southern India: implications for Paleoproterozoic continental reconstruction. *Journal of Earth System Science* 127.
- Belica, M.E., Piispa, E.J., Meert, J.G., Pesonen, L.J., Plado, J., Pandit, M.K., Kamenov, G.D., Celestino, M., 2014. Paleoproterozoic mafic dyke swarms from the Dharwar craton; paleomagnetic poles for India from 2.37-1.88 Ga and rethinking the Columbia supercontinent, *Precambrian Research*, 244, 100-122.
- Bhalla, M. S., Hansraj, A., Prasad Rao, N.T.V., 1980. Paleomagnetic studies of Bangarpet and Sargur dykes of Precambrian age from Karnataka, India. *Geoviews* 8, 181-189.
- Bhowmik, S. K., 2019. The current status of orogenesis in the Central Indian Tectonic Zone: A view from its Southern Margin. *Geological Journal* 54, 2912-2934.
- Bleeker, W., 2003. The late Archean record: a puzzle in ca. 35 pieces. *Lithos* 71, 99-134.
- Butler, R. F., 1992. *Paleomagnetism: magnetic domains to geologic terranes*, Boston: Blackwell Scientific Publications, pp. 248.
- Chadwick, B., Vasudev, V. N., Hegde, G.V., 2000. The Dharwar craton, southern India, interpreted as the result of Late Archaean oblique convergence. *Precambrian Research* 99, 91-111.
- Chardon, D., Jayananda, M., Chetty, T. R., Peucat, J.J., 2008. Precambrian continental strain and shear zone patterns: South Indian case. *Journal of Geophysical Research: Solid Earth* 113.
- Clark, D.A., 1982. Preliminary palaeomagnetic results from the Cuddapah traps of Andhra Pradesh.
- Dash, J. K., Pradhan, S. K., Bhutani, R., Balakrishnan, S., Chandrasekaran, G., Basavaiah, N., 2013. Paleomagnetism of ca. 2.3 Ga mafic dyke swarms in the northeastern Southern Granulite Terrain, India: Constraints on the position and extent of Dharwar craton in the Paleoproterozoic. *Precambrian Research* 228, 164-176.

- Dawson, E.M., Hargraves, R.B., 1994, Paleomagnetism of Precambrian swarms in the Harohalli area, south of Bangalore, India, *Precambrian Research* 69, 157–167.
- Deenen, M. H., Langereis, C. G., van Hinsbergen, D. J., Biggin, A. J. (2011). Geomagnetic secular variation and the statistics of palaeomagnetic directions. *Geophysical Journal International* 186, 509-520.
- Demarest Jr, H. H., 1983. Error analysis for the determination of tectonic rotation from paleomagnetic data. *Journal of Geophysical Research: Solid Earth* 88, 4321-4328.
- Dobmeier, C. J., and Raith, M.M., 2003. Crustal architecture and evolution of the Eastern Ghats Belt and adjacent regions of India. *Geological Society, London, Special Publications* 206, 145-168.
- Evans, D. A. D., and Halls, H. C., 2010. Restoring Proterozoic deformation within the Superior craton. *Precambrian Research* 183, 474-489.
- Fisher, R. A., 1953. Dispersion on a sphere. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 217, 295-305.
- French, J.E., Heaman, L.M., Chacko, T., Srivastava, R.K., 2008. 1891-1883 Ma southern Bastar craton-Cuddapah mafic igneous events, India: a newly recognized large igneous province. *Precambrian Research* 160, 308–322.
- French, J.E., and Heaman, L.M., 2010. Precise U-Pb dating of Paleoproterozoic mafic dyke swarms of the Dharwar craton, India: Implications for the existence of the Neoproterozoic supercraton Sclavia. *Precambrian Research* 183, 416–441.
- Halls, H.C., Kumar, A., Srinivasan, R., Hamilton, M.A., 2007. Paleomagnetism and U-Pb geochronology of easterly trending dykes in the Dharwar craton, India: feldspar clouding, radiating dyke swarms and the position of India at 2.37 Ga, *Precambrian Research* 155, 47-68.
- Hasnain, I., and Qureshy, M. N., 1971. Paleomagnetism and geochemistry of some dikes in Mysore State, India. *Journal of Geophysical Research* 76, 4786-4795.
- Joy, S., Van Der Linde, G., Choudhury, A. K., Deb, G. K., Tappe, S., 2018. Reassembly of the Dharwar and Bastar cratons at ca. 1 Ga: Evidence from multiple tectonothermal events along the Karimnagar granulite belt and Khammam schist belt, southern India. *Journal of Earth System Science* 127, 76.
- Kumar, A., Nagaraju, E., Besse, Jean, Rao, Y.J. Bhaskar, 2012a. New age, geochemical and paleomagnetic data on a 2.21 Ga dyke swarm from south India: Constraints on Paleoproterozoic reconstruction, *Precambrian Research* 220–221, 123–138.
- Kumar, A., Hamilton, M.A., Halls, H.C., 2012b. A Paleoproterozoic giant radiating dyke swarm in the Dharwar craton, southern India, *Geochemistry Geophysics Geosystems* 13, <http://dx.doi.org/10.1029/2011GC003926>.
- Kumar, A., Parashuramulu, V., Nagarju, E., 2015. A 2082 Ma radiating dyke swarm in the Eastern Dharwar craton, southern India and its implications to Cuddapah basin formation, *Precambrian Research* 266, 490-505.

Li, Z. X., and Evans, D. A., 2011. Late Neoproterozoic 40 intraplate rotation within Australia allows for a tighter-fitting and longer-lasting Rodinia. *Geology* 39, 39-42.

Meert, J.G., 2014. In GAD we trust, *Nature Geosciences*, 2, 673-674.

Meert, J.G., Pandit, M.K., Pradhan, V.R., Banks, J.C., Sirianni, R., Stroud, M., Newstead, B., Gifford, J., 2010. The Precambrian tectonic evolution of India: A 3.0 billion year odyssey, *Journal of Asian Earth Sciences* 39, 483-515.

Meert, J.G., Pandit, M.K., Pradhan, V.R., Kamenov, G.D., 2010. Preliminary report on the paleomagnetism of 1.88 Ga dykes from the Bastar and Dharwar cratons, *Gondwana Research* 20, 335-343.

Meert, J.G., Pivarunas, A.K., Miller, S.R., Nutter, R.F., Pandit, M.K., Sinha, A.K. The Precambrian drift history and paleogeography of India, in: Pesonen et al.(eds) *Precambrian Paleomagnetism and Supercontinents*, Elsevier, in press.

Mukherjee, S., Adhikari, A., Vadlamani, R., 2019. Constraining paleoproterozoic (~ 1.7 Ga) collisional orogenesis between the Eastern Dharwar and Bastar cratons: New Sm–Nd garnet isochron and Th–U-total Pb monazite chemical ages from the Bhopalpatnam orogen, central India. *Lithos* 350, 105247.

Myers, J.S., Shaw, R.D., Tyler, I.M., 1996, Tectonic evolution of Proterozoic Australia, *Tectonics* 15, 1431–1446, doi: 10.1029/96TC02356.

Nagaraju, E., Parashuramulu, V., Babu, N.R., Narayana, A.C., 2018a. A 2207 Ma radiating mafic dyke swarm from eastern Dharwar craton, Southern India: Drift history through Paleoproterozoic, *Precambrian Research* 317, 89-100.

Nagaraju, E., Parashuramulu, V., Kumar, A., Sarma, D.S., 2018b. Paleomagnetism and geochronological studies on a 450 km long 2216 Ma dyke from the Dharwar craton, South India, *Physics of the Earth and Planetary Interiors* 274, 222-231.

Piispa, E.J., Smirnov, A.V., Pesonen, L.J., Lindgädevaru, M., Anantha-Murthy, K.S., Devaraju, T.C., 2011. An integrated study of Proterozoic dykes, Dharwar craton, southern India, in Srivastava, R.J. (ed) *Dyke Swarms: Keys for Geodynamic Interpretation*, Springer Publishing, New York, pp. 73-93.

Naqvi, S.M., and Rogers, J.J.W., 1987. *Precambrian Geology of India*. Oxford University Press, Oxford, pp. 223.

Peng, P., Qin, Z., Sun, F., Zhou, X., Guo, J., Zhai, M., Ernst, R.E., 2019. Nature of charnockite and Closepet granite in the Dharwar Craton: Implications for the architecture of the Archean crust. *Precambrian Research* 334, 54-78.

Pivarunas, A.F., Meert, J.G., Pandit, M.K., Sinha, A., 2019. Paleomagnetism and geochronology of mafic dykes from the southern granulite terrane, India: Expanding the Dharwar craton southward, *Tectonophysics* 760, 4-22.

- Pradhan, V.R., Pandit, M.K., Meert, J.G. 2008. A cautionary note on the age of the paleomagnetic pole obtained from the Harohalli dyke swarms, Dharwar craton, southern India, in: Srivastava et al. (eds) *Indian Dykes*, Narosa Publishing House, New Delhi, India, pp. 339-352.
- Prasad, C.V.R.K., Pulla Reddy, V., Subba Rao, K.V., Radhakrishna Murthy, C., 1987. Palaeomagnetism and the crescent shape of the Cuddapah basin, *Geological Society of India Memoirs* 6, 331–347.
- Radhakrishna, T., Joseph, M., 1996. Proterozoic palaeomagnetism of the mafic dyke swarms in the high-grade region of southern India. *Precambrian Research* 76, 31-46.
- Radhakrishna, T., Krishnendu, N., Balasubramonian, G., 2013b. Palaeoproterozoic Indian shield in the global continental assembly: evidence from the palaeomagnetism of mafic dyke swarms, *Earth Science Reviews* 126, 370–389.
- Ramakrishnan, M., and Vaidyanadhan, R., 2008. *Geology of India*. Bangalore, Geological Society of India, 552 pp.
- Santosh, M., Yokoyama, K., Acharyya, S. K., 2004. Geochronology and tectonic evolution of Karimnagar and Bhopalpatnam granulite belts, central India. *Gondwana Research* 7, 501-518.
- Samal, A.K., Srivastava, R., Sinha, L.K., 2015. ArcGIS studies and field relationships of Paleoproterozoic mafic dyke swarms from the south of Devarakonda area, eastern Dharwar craton, southern India: implications for their relative ages, *Journal of Earth Systems Science* 124, 1075-1084.
- Samal, A. K., Srivastava, R. K., Ernst, R. E., Söderlund, U., 2019. Neoproterozoic-mesoproterozoic mafic dyke swarms of the Indian shield mapped using Google Earth™ images and ArcGIS™, and links with large igneous provinces. In *Dyke swarms of the world: A modern perspective*, Springer, Singapore, pp. 335-390.
- Singh, A.P., and Mishra, D.C., 2002. Tectono sedimentary evolution of the Cuddapah basin and Eastern Ghats mobile belt (India) as Proterozoic collision: gravity, seismic and geodynamic constraints. *Journal of Geodynamics* 33, 249–267.
- Singh, A.P., Mishra, D.C., Gupta, S.B., Prabhakar Rao, M.R.K., 2004. Crustal structure and domain tectonics of the Dharwar craton (India): insight from new gravity data. *Journal of Asian Earth Science* 23, 141–152.
- Söderlund, U., Bleeker, W., Demirer, K., Srivastava, R. K., Hamilton, M., Nilsson, M., Srinivas, M., 2019. Emplacement ages of Paleoproterozoic mafic dyke swarms in eastern Dharwar craton, India: Implications for paleoreconstructions and support for a ~ 30° change in dyke trends from south to north. *Precambrian Research* 329, 26-43.
- Srivastava, R. K., Hamilton, M. A., & Jayananda, M., 2011. 2.21 Ga large igneous province in the Dharwar Craton, India. In *International Symposium on Large Igneous Provinces of Asia, Mantle Plumes and Metallogeny, Irkutsk, Russia, Extended Abstract* (pp. 263-266).

Stein, H.J., Hannah, J.L., Zimmerman, A., Markey, R. Sarkar, S.C., Pal, A.B., 2004. A 2.5 Ga porphyry Cu-Mo-Au deposit at Malanjhand, central India: Implications for Late Archean continental assembly, *Precambrian Research* 134, 189-226.

Stein, H., Hannah, J., Zimmerman, A. & Markey, R., 2006. Mineralization and deformation of the Malanjhand terrane (2,490–2,440 Ma) along the southern margin of the Central Indian Tectonic Zone. *Mineralium Deposita* 40, 755–765.


Tauxe, L. Shaar, R., Jonestrask, L., Swanson-Hysell, N.L., Minnett, R., Koppers, A.A.P., Constable, G.C., Jarboe, N., Gaastra, K. Fairchild, L. 2016. PmagPy: Software package for paleomagnetic data analysis and a bridge to the Magnetism Information Consortium (MagIC) Database, *Geochemistry, Geophysics Geosystems* 17, doi:10.1002/2016GC006307.



Tauxe, L., and Kent, D.V., 2004. A simplified statistical model for the geomagnetic field and the detection of shallow bias in paleomagnetic inclinations: was the ancient magnetic field dipolar? In *Timescales of the Paleomagnetic Field*, Blackwell Publishing Ltd., pp. 101-115.

Valet, J., Besse, J., Kumar, A., Vadaque-Chanat, S., Philippe, E., 2014. The intensity of the geomagnetic field from 2.4 Ga old Indian dykes, *Geochemistry, Geophysics, Geosystems* 15, 2426-2437.



Vanderkluysen, L., Mahoney, J. J., Hooper, P. R., Sheth, H. C., and Ray, R., 2011. The feeder system of the Deccan Traps (India): insights from dike geochemistry. *Journal of Petrology* 52, 315-343.





Venkatesh, A. S., Rao, G. P., Rao, N. P., Bhalla, M.S., 1987. Palaeomagnetic and geochemical studies on dolerite dykes from Tamil Nadu, India. *Precambrian Research* 34, 291-310.







- △ North mean
- north sites
- South mean
- south sites







Declaration of interests

□ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Anthony Pivarunas and Joseph Meert